

Letter to Editor

Open letter to journal editors on: international consensus radiochemistry nomenclature guidelines

Heinz H Coenen¹, Antony D Gee², Michael Adam³, Gunnar Antoni⁴, Cathy S Cutler⁵, Yasuhisa Fujibayashi⁶, Jae Min Jeong⁷, Robert H Mach⁸, Thomas L Mindt⁹, Victor W Pike¹⁰, Albert D Windhorst¹¹

¹Research Centre Jülich, Jülich, Germany; ²King's College London, London, UK; ³TRIUMF, Vancouver, Canada; ⁴Uppsala University, Uppsala, Sweden; ⁵Brookhaven National Laboratory, Upton, USA; ⁶Keio University, Tokyo, Japan; ⁷Seoul National University, Seoul, South Korea; ⁸University of Pennsylvania, Philadelphia, USA; ⁹Ludwig Boltzmann Institute Applied Diagnostics, Vienna, Austria; ¹⁰National Institute of Mental Health, Bethesda, USA; ¹¹VU University Medical Center, Amsterdam, The Netherlands

Received January 10, 2018; Accepted January 10, 2018; Epub February 5, 2018; Published February 15, 2018

International consensus nomenclature guidelines-summary [1]

Measures of radioactivity

Unit of activity

“Radioactivity” is a physical *phenomenon*, defined as the *property* of certain nuclei to spontaneously fragment or rearrange, resulting in the emission of radiation.

“Activity” is the *quantitative measure* of radioactivity: The number of nuclear decays, occurring in a given quantity of material over a certain time interval, divided by that time interval.

“Becquerel” (Bq) is the agreed SI derived unit for the *quantity* of activity.

Pre-SI units (e.g. imperial units) (e.g. mCi, Ci) can also be used, but must be placed in parentheses after the stated SI units.

N.B.: The correct terms “radioactive” and “non-radioactive” must not be replaced by the lab-jargon “hot” and “cold”, respectively, in public or official documents.

Molar activity (A_m) and specific activity (A_s)

Molar activity is the measured activity per mole of compound; measured in Bq/mol (GBq/ μ mol).

Specific activity is the measured activity per gram of compound; measured in Bq/g (GBq/ μ g).

Due to radioactive decay, the instant of time of measurement must be stated; for example: “The molar activity was 50 GBq/ μ mol 2 h after the end of synthesis”.

Apparent molar activity and apparent specific activity: The terms apparent molar activity and apparent specific activity take into account the amounts of the labelled and non-radiolabelled impurities present (using moles, or weight, respectively).

Effective molar activity and effective specific activity: The terms “effective molar activity” and “effective specific activity” address the chemically, biologically or pharmacologically ‘active’ fraction of radioactive and non-radioactive materials in a sample, competing with the labelled product in its chemical or biological reactions. In this case, the “effectivity” must be determined by an additional analytical process; e.g. receptor or enzyme binding assay, side-product analysis, etc.

Other terms like “pseudo-specific activity” etc. must not be used.

‘No-carrier-added’, ‘carrier added’ and ‘carrier-free’: The non-quantitative terms, “no-carrier-added” (n.c.a.) and “carrier-added” (c.a.) can be used as a practical, qualitative indication of specific or molar activity.

The term “carrier-free” (c.f.) can only be used in the rare case where the theoretical maximum

Radiochemistry Nomenclature

specific or molar activity is unambiguously proven.

Measures involving determination of amounts of material (e.g. molar activity or radiochemical purity), should be accompanied by a clear description of the method of detection.

Radionuclide and radioisotope descriptors

The enrichment of a chemical compound with an isotope (stable or radioactive) of one or more of the elements, of which it is constituted, is indicated by the symbol of the element (E) together with its mass number (A) (as a superscript in front) within square brackets, [^AE], immediately preceding the compound's name or chemical formula.

Example: [²H,¹⁴C]benzene, or [²H,¹⁴C]C₆H₆, represent the compound *benzene*, enriched or labelled, with stable deuterium and radioactive carbon-14, respectively.

For rules designating labelling positions (e.g. L-[*methyl*-¹⁴C] methionine or L-[*carboxyl*-¹⁴C] methionine) see: <http://goldbook.iupac.org/pages/about.html>.

These rules apply equally to organic, inorganic and organometallic compounds and complexes: e.g. [²²³Ra]RaCl₂, [^{99m}Tc]NaTcO₄, [^{99m}Tc]Tc-MDP, [⁶⁸Ga]Ga-DOTA-TATE, etc.

As square brackets are also used to denote metal complexes, care should be taken to avoid confusion with radionuclide descriptors, e.g. [^{99m}Tc][Tc(CO)₃(OH₂)₃]⁺, [¹¹¹In][In(DTPA)]²⁺, etc.

N.B.: The symbol for isotopic enrichment [^AE] should be treated like a syllable, and thus only be hyphenated at the end of a line of text.

Conversely, do not use nuclide symbols in square brackets in combination with nouns and verbs.

Examples of the use of square brackets and hyphens:

Correct	Incorrect
L-[¹³ N]alanine, (S)-[¹³ N]alanine	[¹³ N]L-alanine, (S)- ¹³ N-alanine, L-[¹³ N]-alanine
[¹⁸ F]fluorobenzene	[¹⁸ F]benzene (no fluorine atom in benzene)

[¹⁸ F]fluorobenzene	[^{99m} Tc]DTPA ²⁻ , ^{99m} Tc-DTPA ²⁻ (no Tc in chelator)
¹¹ C-compound, ¹²⁵ I-substitution,	[¹¹ C]compound, [¹²⁵ I]-substitution,
¹⁸ F-derivative, ⁶⁸ Ga-conjugate chemical	[¹⁸ F]-derivative, [⁶⁸ Ga]conjugate (no compound names)
⁷⁷ Br-reagent	
¹¹ C -labelling, ⁶⁴ Cu-labelling,	[¹¹ C]labelling, [⁶⁴ Cu]-labelling,
¹⁸ F-fluorination (with hyphen!)	[¹⁸ F]-(radio)-fluorination (no chemical compound names)

The terms '(radio) isotope' and '(radio) nuclide' are often used incorrectly in texts, inferring "isotope" means "radioactive nuclide" or even "labelled compound".

'Nuclide' indicates an atom, characterised by its numbers of protons (identifying its elemental nature) and of nucleons (indicating its mass).

'Isotopes' are nuclides of the same element (same proton number), but having different numbers of neutrons (hence different atomic mass).

Isotopic nuclides of different energy state are called isomeric nuclides, isomeric isotopes or 'isomers', such as technetium -99 g and -99 m.

N.B.: All (radio) isotopes are (radio) nuclides, while the reverse is not true!

Radiochemical yield (RCY)

The "radiochemical yield" is the amount of activity in the product expressed as the percentage (%) of related starting activity utilized in the considered process (e.g. synthesis, separation, etc.). The quantity of both must relate to the same radionuclide and be decay corrected to the same point in time before the calculation is made.

(Reported measures of RCY should indicate, whether a product was isolated or not).

Colloquial expressions for "radio-yield" found in literature, e.g. 'radiochemical conversion', 'analytical radiochemical yield', 'radio-HPLC yield' must not be used as a surrogate for 'radiochemical yield' or 'radiochemical purity', respectively.

Radiochemistry Nomenclature

The following text may serve as examples for good practice, when describing a radiochemical yield:

“The radiochemical yield was 67% (based on HPLC analysis of the crude product)”.

“The radiochemical yield* of “Y” was 67%”, with the following as a footnote:

* “determined by radio-HPLC analysis of the crude product”, or * “non-isolated, estimated by radio-HPLC”, or, in the general experimental section: “All radiochemical yields were determined by HPLC of the crude product, unless stated otherwise”, or alternatively use: “The radiochemical purity of the crude product was 67%”, or: “The radiochemical yield of “Y” determined from an aliquot of the reaction solution amounted to 67%”, or: “The radiochemical yield of crude “Y” was 67% based on the amount of activity eluted from the HPLC column”.

Expressions such as ‘conversion’ or ‘incorporation’, however, may be used in a semantic sense and even be indispensable in context of mechanistic discussions. For example, The ‘conversion’ (or ‘incorporation’) proceeded with a 50% yield. Here it is clear from the context, that the radiochemical yield of the conversion is intended.

Activity yield

The “activity yield” is the amount of radioactive product expressed in Bq (MBq, GBq), which is obtained from a starting amount of activity (e.g. produced at a cyclotron) and is not corrected for decay.

This term is useful, or necessary to indicate the efficiency of a labelling procedure. The activity yield is, of course, dependent on the effectiveness and duration of all technical manipulations used, in addition to the yield of the labelling reaction.

Address correspondence to: Heinz H Coenen, Research Centre Jülich, Jülich, Germany. E-mail: h.h.coenen@fz-juelich.de; Antony D Gee, King’s College London, London, UK. E-mail: antony.gee@kcl.ac.uk

Reference

- [1] Coenen HH, Gee AD, Adam M, Antoni G, Cutler CS, Fujibayashi Y, Jeong JM, Mach RH, Mindt TL, Pike VW, Windhorst AD. Consensus nomenclature rules for radiopharmaceutical chemistry-setting the record straight. *Nucl Med Biol* 2017; 55: v-xi.